

1

2

3

4

<sup>1</sup> Department of Earth and Environmental Engineering, Columbia University, New York, NY.

<sup>2</sup> Department of Earth System Science, Stanford University, Stanford, CA.

<sup>3</sup> Department of Civil and Environmental Engineering, Massachusetts Institute of Technology, Cambridge, MA.

<sup>4</sup> Department of Global Ecology, Carnegie Institution of Washington, Stanford, CA.

<sup>5</sup> Department of Earth, Atmospheric and Planetary Sciences, Massachusetts Institute of Technology, Cambridge, MA.

<sup>6</sup> University Space Research Association, Columbia, MD.

<sup>7</sup> Global Modeling and Assimilation Office, NASA Goddard Space Flight Center, Greenbelt, MD.

<sup>8</sup> Department of Earth, Environment and Planetary Sciences, Brown University, Providence, RI.

<sup>9</sup> The Earth Institute, Columbia University, New York, NY.

\*Correspondence to: jg3405@columbia.edu.

5           **The terrestrial biosphere and atmosphere interact through a series of feedback**  
6 **loops. Variability in terrestrial vegetation growth and phenology can modulate fluxes of**  
7 **water and energy to the atmosphere, and thus affect the climatic conditions that in turn**  
8 **regulate vegetation dynamics. Here we analyze satellite observations of solar-induced**  
9 **fluorescence, precipitation, and radiation using a multivariate statistical technique. We**  
10 **find that biosphere-atmosphere feedbacks are globally widespread and regionally**  
11 **strong: they explain up to 30% of precipitation and surface radiation variance.**  
12 **Substantial biosphere-precipitation feedbacks are often found in regions that are**  
13 **transitional between energy and water limitation, such as semi-arid or monsoonal**  
14 **regions. Substantial biosphere-radiation feedbacks are often present in several**  
15 **moderately wet regions and in the Mediterranean, where precipitation and radiation**  
16 **increase vegetation growth. Enhancement of latent and sensible heat transfer from**  
17 **vegetation accompanies this growth, which increases boundary layer height and**  
18 **convection, affecting cloudiness, and consequently incident surface radiation. Enhanced**  
19 **evapotranspiration can increase moist convection, leading to increased precipitation.**  
20 **Earth system models underestimate these precipitation and radiation feedbacks mainly**  
21 **because they underestimate the biosphere response to radiation and water availability.**  
22 **We conclude that biosphere-atmosphere feedbacks cluster in specific climatic regions**  
23 **that help determine the net CO<sub>2</sub> balance of the biosphere.**

24           By influencing the partitioning of turbulent fluxes at the surface<sup>1</sup>, soil moisture and  
25 temperature can affect climatic variability<sup>2</sup>. Biospheric variability, in terms of both  
26 phenology and stomatal regulation, also strongly modulates turbulent fluxes of both water  
27 and energy<sup>3</sup>. Since biospheric variability is regulated by vegetation phenology and root zone  
28 soil moisture, it exhibits longer (e.g. multi-month) memory compared to the more commonly  
29 studied surface soil moisture and temperature state. Therefore, an understanding of

30 biosphere-atmosphere interactions has the potential to improve seasonal to interannual  
31 climatic predictions<sup>4,5,6</sup>, and improve predictions of vegetation resilience to climate  
32 anomalies<sup>7</sup>. However, global variations in the strength of biosphere-atmosphere feedbacks  
33 remain unknown, in part because of the difficulty of observing biospheric fluxes<sup>8</sup>.

34       Recent advancements in space-borne observations of solar-induced fluorescence (SIF)  
35 have enabled for the first-time a global proxy for gross primary productivity (GPP) and  
36 vegetation phenology. SIF is a by-product of photosynthesis<sup>9</sup> related to light-use efficiency  
37 (LUE) and to the fraction of absorbed photosynthetic active radiation (fAPAR)<sup>10</sup>. On a  
38 canopy or regional scale and at a monthly resolution it is nearly proportional to GPP across  
39 various ecosystems. This large-scale correspondence is strongly related to the changes in  
40 canopy structure and phenology on absorbed photosynthetic active radiation, in addition to  
41 the more subtle changes in LUE<sup>11,12,13,14</sup>. SIF is also generally highly correlated with  
42 evapotranspiration (ET)<sup>15</sup> (Supplementary Fig. 1) and correlates with vegetation-driven  
43 changes in surface albedo. Here, we use SIF as an integrated measure of vegetation  
44 variability, capturing both growth and changes in photosynthetic capacity (Methods).

45       Previous studies of land-atmosphere interactions have typically relied on correlations  
46 between land and atmospheric variables<sup>16,17,18</sup>. However, these variables seasonally coevolve,  
47 and thus it is difficult to determine whether one variable is causally forcing the other, or if the  
48 two are both driven by separate factors<sup>19,20</sup>. Here, these shortcomings are overcome by  
49 employing a Multivariate Conditional Granger causality (MVGC) statistical technique using  
50 vector autoregressive models (VARs)<sup>21</sup>. This method determines both the strength of the  
51 predictive mechanism between variables and the time scale over which these links occur  
52 (Methods).

## 53 **MVGC observational data forcings**

54       We apply the MVGC VAR statistical technique to eight years of monthly SIF  
55 measurements from the Global Ozone Monitoring Experiment 2 (GOME-2) sensor<sup>22</sup>. SIF-  
56 precipitation interactions are assessed using remote sensing-based estimates from the Global  
57 Precipitation Climatology Project (GPCP)<sup>23</sup> and SIF-radiation interactions are assessed using  
58 photosynthetic active radiation (PAR) from Clouds and the Earth's Radiant Energy System  
59 (CERES)<sup>24</sup>. We also use surface air temperature reanalysis data from ERA-Interim<sup>25</sup>, as  
60 temperature can independently impact and interact with photosynthetic activity<sup>18</sup>. SIF data is  
61 relatively noisy, and thus spatial averaging is used to smooth it prior to analysis (Methods). It  
62 should be acknowledged that the smoothing could distort results in highly heterogeneous  
63 regions where signals from various biomes may be aggregated. Note that, although the linear  
64 scaling factor between monthly SIF and GPP varies between ecosystems and climates<sup>12</sup> the  
65 pixel-by-pixel data normalization used here removes the geographical variations of this factor  
66 (Methods). The analysis presented here is independent of the scaling factor.

67       To identify biosphere-atmosphere coupled feedbacks, we first examine their  
68 directional sub-components, i.e. the *atmospheric forcing* (atmosphere → biosphere), as  
69 assessed by the response of SIF (GPP) to atmospheric drivers (the fraction of variance in SIF  
70 explained by precipitation and PAR), and the *biospheric forcing* (biosphere → atmosphere),  
71 as assessed by precipitation and PAR response to SIF (the fraction of variance in  
72 precipitation and PAR explained by SIF) (Fig. 1). An F-test with a null-hypothesis of 0-  
73 Granger causality (G-causality) (p-value <0.1) is used. The total feedback strength is then  
74 defined as the product of these two directional components (Fig. 2). The sign of the feedback  
75 is defined as the sign of the first order coefficient of the VAR model from the G-causality  
76 analysis. To ensure the results presented here are robust and independent of the seasonal  
77 cycle (i.e. due to land-atmosphere interactions), a bootstrap test that conserves the seasonal

78 cycle but breaks the causality by shuffling months from different years is used  
79 (Supplementary Fig. 2) and clearly destroys the feedback.

80 Globally, precipitation positively explains the highest fraction of biosphere (SIF)  
81 variability in regions that are transitional between wet and dry climates, e.g. semi-arid or  
82 monsoonal (Fig. 1a), consistent with previous studies<sup>7,16</sup>. Many of these regions also have  
83 high fractions of C4 plants<sup>26</sup>, which have higher water use efficiency than C3 species<sup>27</sup>, and  
84 are therefore expected to be more sensitive to water limitations. The impact of the biosphere  
85 on precipitation (Fig. 1b), as assessed by the G-causality of SIF on precipitation, is seen in  
86 seasonally dry regions where increases in GPP, in response to increased soil moisture and  
87 vegetation growth, is linked with higher latent heat flux and reduced sensible heat flux  
88 (Supplementary Fig. 1). Although the impact of SIF on precipitation is less widespread than  
89 that of precipitation on SIF, it is significant in many of the same regions. The feedbacks are  
90 almost always positive because the monthly positive effect of evapotranspiration on moist  
91 convection dominates negative feedback pathways induced by mesoscale surface  
92 heterogeneity<sup>28</sup> and the effects of changing albedo. The time scales involved in the feedback  
93 mechanisms can vary between regions. The subseasonal signal may represent variability due  
94 to early greening induced by increased water supply or to browning induced by water stress,  
95 while seasonal and interannual signals may indicate changes in vegetation growth regulated  
96 by water availability during cell division. The strongest signals are detected subseasonally in  
97 monsoonal Australia, seasonally in Eastern Asia, and both seasonally and interannually in the  
98 Sahel and Southern African Monsoonal regions (Supplementary Fig. 3). The dominance of  
99 seasonal and interannual time scales in the Sahel, related to biomass variability, is consistent  
100 with previous understanding<sup>6,29</sup>.

101 PAR has the greatest impact on biosphere fluxes (Fig. 1c) in regions where  
102 photosynthesis and vegetation growth is energy limited such as the high latitudes, humid

regions of the Eastern US, parts of the Mediterranean, and tropical rainforest regions<sup>30,31</sup>. This agrees with the findings of previous studies showing that net primary production (NPP) in these regions is driven by radiation<sup>18</sup>. The biosphere exerts control on PAR in the Eastern US, central Eurasia, African deciduous woodlands as well as in the European Mediterranean region (Fig. 1d). In these very dry or very wet regions, ecosystems rarely enter the transitional regime where stomatal closure depends on soil moisture, and increases in SIF are accompanied by increases in both sensible and latent heat (Supplementary Fig. 1)<sup>32</sup>. The increased sensible heat flux leads to a deeper boundary layer and reduced cloud cover (Supplementary Fig. 4), therefore increasing PAR (Fig. 1d). In the Eastern US, the increase in PAR is mostly attributed to a reduction of low- and mid-level (i.e. congestus) cumulus clouds, typical of summer conditions in this humid climate (Supplementary Fig. 4). By contrast, in the European Mediterranean, PAR is most sensitive to mid- and high-level clouds. In central Eurasia all cloud cover levels negatively impact surface PAR but high-level clouds are the primary reason for the PAR change. The strongest feedbacks between SIF and PAR tend to be on a seasonal scale indicating an increase in ecosystem-scale photosynthetic capacity due to vegetation growth, with exceptions in Madagascar, Australia and central Eurasia where subseasonal and interannual feedbacks dominate (Supplementary Fig. 3). In all PAR feedback regions, PAR is also negatively correlated with precipitation (Supplementary Fig. 4). We note that the European Mediterranean has been highlighted as a hotspot of land-atmosphere coupling in an earlier modeling study, emphasizing the strong coupling between surface turbulent fluxes and the boundary layer response in the region<sup>33</sup>. While a similar coupling mechanism may occur in other regions, they do not exhibit a strong response because other processes (e.g. topography, different land-ocean circulation...) overshadow the regional impact of the biosphere there.

## MVGC observational data coupled feedbacks

The results of the *atmospheric* and *biospheric forcings* (Fig. 1) are combined to determine the total variance explained in the coupled biosphere-atmosphere system (Fig. 2 and Supplementary Fig. 5). Hotspot regions for the precipitation → SIF → precipitation feedback (Fig. 2a) - which can explain up to 20-30% of the observed precipitation variance - are concentrated in grasslands and savannas (transitional zones) such as monsoonal regions in the Sahel, Eastern India and Northern Australia, as well as the African savanna, Madagascar and the Brazilian savannas. There are other monsoonal regions that despite large shifts in rainfall during the year are not hotspots either due to a lack of ET response to precipitation<sup>34</sup>, or a lack of precipitation response to changes in ET<sup>35</sup>. An example of this is the Central Great Plains in North America (a hotspot per previous modeling-based studies of soil moisture-atmosphere interactions<sup>36</sup>), where soil moisture has been shown to have a weak triggering effect on precipitation<sup>20,37</sup>. Indeed, summertime precipitation in this area is dominated by eastward propagating mesoscale convective systems mostly independent of the land surface<sup>38</sup>.

The PAR → SIF → PAR feedback (Fig. 2b) has hotspots (20-30% of explained variance) in the humid Eastern United States, Southern Brazil, as well as in the Mediterranean basin in Europe. By contrast, in the tropical rainforest regions of Africa and South America there is little response detected for the full feedback loops with either precipitation or PAR (Fig. 2 and Supplementary Fig. 5) suggesting that other factors (such as ecosystem characteristics<sup>39</sup>) dominate the variability of the biosphere there.

Although feedbacks between the biosphere and atmosphere are detected in almost all regions, several 'hotspot families' stand out: 1) regions that are either semi-arid or monsoonal for the precipitation feedback and 2) humid regions (the Eastern US) and the Mediterranean

for the PAR feedback. No regions exhibit both feedback pathways; one always dominates the other when it is present.

### **MVGC ESM analysis**

The distribution of feedbacks in the observational record is next used to assess Earth System Models (ESMs) (Supplementary Table 1). The distributions of feedback strengths for model and observational results (Fig. 3) summarize the differences between the biosphere-atmosphere feedback detected by each CMIP5 model (Supplementary Figs 6, 7 and 8) and the observational record. In the model analysis, GPP is used as a proxy for the biosphere in lieu of SIF. Our results are normalized in terms of explained variance for each pixel so that the proportionality factor of SIF and GPP does not impact the pixel-wise metric results. To increase robustness, 50 years of data are used for the model analysis (1956-2005) rather than the shorter period we are constrained by for the observational analysis<sup>40</sup>.

The median of all ESMs fall below the first quartile of the observational data results for the precipitation → biosphere → precipitation feedback (Fig. 3a). Models significantly underestimate the magnitude and the range of both the *atmospheric* and *biospheric forcings* (except for CMCC-CESM) (Supplementary Fig. 6), although underestimation is more severe in the case of the precipitation → biosphere component. The observational PAR → biosphere → PAR feedback strength (Fig. 3b) also has a higher median value than that of the ESMs. Both the precipitation and PAR *atmospheric forcings* are underestimated because of photosynthesis misrepresentation in ESMs (Supplementary Fig. 6)<sup>41</sup>. Despite some spatial similarities between modeled feedbacks and observational results (Supplementary Figs 7 and 8), models systematically underestimate the impact of the biosphere on precipitation, and noticeably miss the variance explained by observations in monsoonal Australia. On the other hand, the modeled impact of the biosphere on PAR varies drastically between models and can

be either over- or under-estimated (Supplementary Fig. 6). These inter-model discrepancies are likely due to the misrepresentation of convection in models, and the challenges of correctly representing it over land regions<sup>42,43</sup>. Interestingly, in general, ESM errors in representing the atmospheric forcing on the biosphere are even more severe than errors in representing the biospheric forcing on the atmosphere. This suggests that better representations of photosynthesis and water stress sensitivities would have a larger impact on improving the ESM representation of biosphere-atmosphere feedbacks, than improved convection representation.

This study provides the first causal observational diagnostic of biosphere-atmosphere feedbacks on subseasonal to interannual time scales. These feedbacks are strong in semi-arid and monsoonal regions, which are key in determining whether the yearly global terrestrial biosphere acts as a net CO<sub>2</sub> source or sink<sup>7,16</sup>. As such biosphere-atmosphere feedbacks regulate interannual hydrology and climate in these regions as well as the global carbon cycle. Additionally, due to the high percentages of atmospheric variability explained by vegetation processes, subseasonal and seasonal climate predictions can greatly benefit from better vegetation characterization in ESMs. In turn this will improve subseasonal to seasonal climate and hydrologic forecasts, which are crucial for optimizing management decisions pertaining to food security, water supplies, and disaster management such as droughts and heat waves.

## 194 Main Text References

- 195 1. Bateni, S. M. & Entekhabi, D. Relative efficiency of land surface energy balance  
196 components. *Water Resour. Res.* **48**, 1–8 (2012).
- 197 2. Koster, R. D., Suarez, M. J. & Heiser, M. Variance and Predictability of Precipitation  
198 at Seasonal-to-Interannual Timescales. *J. Hydrometeorol.* **1**, 26–46 (2000).
- 199 3. van den Hurk, B. J. J. M., Viterbo, P. & Los, S. O. Impact of leaf area index  
200 seasonality on the annual land surface evaporation in a global circulation model. *J.*  
201 *Geophys. Res.* **108**, 5.1-5.7 (2003).
- 202 4. Guo, Z., Dirmeyer, P. A., Delsole, T. & Koster, R. D. Rebound in atmospheric  
203 predictability and the role of the land surface. *J. Clim.* **25**, 4744–4749 (2012).
- 204 5. Koster, R. D. *et al.* The Second Phase of the Global Land–Atmosphere Coupling  
205 Experiment: Soil Moisture Contributions to Subseasonal Forecast Skill. *J.*  
206 *Hydrometeorol.* **12**, 805–822 (2011).
- 207 6. Zeng, N., Neelin, J., Lau, K. & Tucker, C. Enhancement of Interdecadal Climate  
208 Variability in the Sahel by Vegetation Interaction. *Science* **286**, 1537–1540 (1999).
- 209 7. Poulter, B. *et al.* Contribution of semi-arid ecosystems to interannual variability of the  
210 global carbon cycle. *Nature* **509**, 600–603 (2014).
- 211 8. Koster, R. D. *et al.* On the nature of soil moisture in land surface models. *J. Clim.* **22**,  
212 4322–4335 (2009).
- 213 9. Porcar-Castell, A. *et al.* Linking chlorophyll a fluorescence to photosynthesis for  
214 remote sensing applications: Mechanisms and challenges. *J. Exp. Bot.* **65**, 4065–4095  
215 (2014).

- 216 10. Guanter, L. *et al.* Global and time-resolved monitoring of crop photosynthesis with  
217 chlorophyll fluorescence. *Proc. Natl. Acad. Sci. U. S. A.* **111**, E1327-33 (2014).
- 218 11. Zhang, Y. *et al.* Remote Sensing of Environment Consistency between sun-induced  
219 chlorophyll fluorescence and gross primary production of vegetation in North  
220 America. *Remote Sens. Environ.* **183**, 154–169 (2016).
- 221 12. Frankenberg, C. *et al.* New global observations of the terrestrial carbon cycle from  
222 GOSAT: Patterns of plant fluorescence with gross primary productivity. *Geophys. Res.*  
223 *Lett.* **38**, L17706 (2011).
- 224 13. Frankenberg, C., O'Dell, C., Guanter, L. & McDuffie, J. Remote sensing of near-  
225 infrared chlorophyll fluorescence from space in scattering atmospheres: Implications  
226 for its retrieval and interferences with atmospheric CO<sub>2</sub> retrievals. *Atmos. Meas.*  
227 *Tech.* **5**, 2081–2094 (2012).
- 228 14. Wood, J. D. *et al.* Multi-scale analyses reveal robust relationships between gross  
229 primary production and solar induced fluorescence. *Geophys. Res. Lett.* **In Review**,  
230 533–541 (2016).
- 231 15. Schlesinger, W. H. & Jasechko, S. Transpiration in the global water cycle. *Agric. For.*  
232 *Meteorol.* **189–190**, 115–117 (2014).
- 233 16. Ahlström, A. *et al.* The dominant role of semi-arid ecosystems in the trend and  
234 variability of the land CO<sub>2</sub> sink. *Science (80-. ).* **348**, 895–899 (2015).
- 235 17. Beer, C. *et al.* Terrestrial gross carbon dioxide uptake: global distribution and  
236 covariation with climate. *Science* **329**, 834–838 (2010).
- 237 18. Nemani, R. R. *et al.* Climate-driven increases in global terrestrial net primary  
238 production from 1982 to 1999. *Science* **300**, 1560–1563 (2003).

- 239 19. Sugihara, G. *et al.* Detecting causality in complex ecosystems. *Science* **338**, 496–500  
240 (2012).
- 241 20. Tuttle, S. & Salvucci, G. Empirical evidence of contrasting soil moisture–precipitation  
242 feedbacks across the United States. **352**, 825–828 (2016).
- 243 21. Barnett, L. & Seth, A. K. The MVGC multivariate Granger causality toolbox: A new  
244 approach to Granger-causal inference. *J. Neurosci. Methods* **223**, 50–68 (2014).
- 245 22. Joiner, J. *et al.* Global monitoring of terrestrial chlorophyll fluorescence from  
246 moderate spectral resolution near-infrared satellite measurements: methodology,  
247 simulations, and application to GOME-2. *Atmos. Meas. Tech. Discuss.* **6**, 3883–3930  
248 (2013).
- 249 23. Pendergrass, A. & N. C. for A. R. S. (Eds). The Climate Data Guide: GPCP  
250 (Monthly): Global Precipitation Climatology Project. 1–2 (2016). Available at:  
251 [https://climatedataguide.ucar.edu/climate-data/gpcp-monthly-global-precipitation-](https://climatedataguide.ucar.edu/climate-data/gpcp-monthly-global-precipitation-climatology-project)  
252 [climatology-project](https://climatedataguide.ucar.edu/climate-data/gpcp-monthly-global-precipitation-climatology-project).
- 253 24. Wielicki, B. A. *et al.* Clouds and the Earth’s Radiant Energy System (CERES): An  
254 Earth Observing System Experiment. *Bull. Amer. Meteor. Soc.* **77**, 853–868 (1996).
- 255 25. Dee, D. P. *et al.* The ERA-Interim reanalysis: Configuration and performance of the  
256 data assimilation system. *Q. J. R. Meteorol. Soc.* **137**, 553–597 (2011).
- 257 26. Still, C. J., Berry, J. A., Collatz, G. J. & DeFries, R. S. Global distribution of C 3 and  
258 C 4 vegetation: Carbon cycle implications. *Global Biogeochem. Cycles* **17**, 6-1-6–14  
259 (2003).
- 260 27. Ghannoum, O. C4 photosynthesis and water stress. *Ann. Bot.* **103**, 635–644 (2009).
- 261 28. Guillod, Benoit P. (Institute for Atmospheric and Climate Science, Department of

- 262 Environmental Systems Science, E. Z., Orlowsky, B., Miralles, D. G., Teuling, A. J. &  
263 Seneviratne, S. I. Reconciling spatial and temporal soil moisture effects on afternoon  
264 rainfall. *Nat. Commun.* **6**, 6443 (2015).
- 265 29. Charney, J. G. Dynamics of deserts and drought in the Sahel. *Q. J. R. Meteorol. Soc.*  
266 **101**, 193–202 (1975).
- 267 30. Anber, U., Gentine, P., Wang, S. & Sobel, A. H. Fog and rain in the Amazon. *Proc.*  
268 *Natl. Acad. Sci.* **112**, 11473–11477 (2015).
- 269 31. Brando, P. M. *et al.* Seasonal and interannual variability of climate and vegetation  
270 indices across the Amazon. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 14685–90 (2010).
- 271 32. Seneviratne, S. I. *et al.* Investigating soil moisture-climate interactions in a changing  
272 climate: A review. *Earth-Science Rev.* **99**, 125–161 (2010).
- 273 33. Seneviratne, S. I., Lüthi, D., Litschi, M. & Schär, C. Land-atmosphere coupling and  
274 climate change in Europe. *Nature* **443**, 205–209 (2006).
- 275 34. Dirmeyer, P. A. The terrestrial segment of soil moisture-climate coupling. *Geophys.*  
276 *Res. Lett.* **38**, L16702 (2011).
- 277 35. Koster, R. D. & Suarez, M. J. Impact of Land Surface Initialization on Seasonal  
278 Precipitation and Temperature Prediction. *J. Hydrometeorol.* **4**, 408–423 (2003).
- 279 36. Koster, R. D. *et al.* GLACE: The Global Land – Atmosphere Coupling Experiment.  
280 Part I: Overview. *J. Hydrometeorol.* **7**, 611–625 (2006).
- 281 37. Findell, K. L., Gentine, P., Lintner, B. R. & Kerr, C. Probability of afternoon  
282 precipitation in eastern United States and Mexico enhanced by high evaporation. *Nat.*  
283 *Geosci.* **4**, 434–439 (2011).
- 284 38. Storer, R. L., Zhang, G. J. & Song, X. Effects of convective microphysics

285 parameterization on large-scale cloud hydrological cycle and radiative budget in  
 286 tropical and midlatitude convective regions. *J. Clim.* **28**, 9277–9297 (2015).

287 39. Levine, N. M. *et al.* Ecosystem heterogeneity determines the ecological resilience of  
 288 the Amazon to climate change. *Proc. Natl. Acad. Sci.* (2015).  
 289 doi:10.1073/pnas.1511344112

290 40. Findell, K. L., Gentine, P., Lintner, B. R. & Guillod, B. P. Data Length Requirements  
 291 for Observational Estimates of Land–Atmosphere Coupling Strength. *J.*  
 292 *Hydrometeorol.* **16**, 1615–1635 (2015).

293 41. Zhou, S., Duursma, R. A., Medlyn, B. E., Kelly, J. W. G. & Prentice, I. C. How should  
 294 we model plant responses to drought? An analysis of stomatal and non-stomatal  
 295 responses to water stress. *Agric. For. Meteorol.* **182–183**, 204–214 (2013).

296 42. Bony, S. *et al.* Clouds, circulation and climate sensitivity. *Nat. Geosci.* **8**, 261–268  
 297 (2015).

298 43. Zhao, M. *et al.* Uncertainty in model climate sensitivity traced to representations of  
 299 cumulus precipitation microphysics. *J. Clim.* **29**, 543–560 (2016).

300

## **Corresponding author**

Correspondence and requests for materials should be addressed to JKG (jg3405@columbia.edu).

## **Acknowledgements**

The authors would like to thank Guido Salvucci and Upmanu Lall for discussion on the Granger causality, Randal Koster for initial discussion of the paper, and Joanna Joiner for providing GOME-2 data. This project was supported by both a NASA Earth Science and Space Fellowship as well as a DOE GOAmazon grant.

“We acknowledge the World Climate Research Programme's Working Group on Coupled Modelling, which is responsible for CMIP, and we thank the climate modeling groups (listed in Table S1 of this paper) for producing and making available their model output. For CMIP the U.S. Department of Energy's Program for Climate Model Diagnosis and Intercomparison provides coordinating support and led development of software infrastructure in partnership with the Global Organization for Earth System Science Portals.”

## **Author Contributions**

JKG, AGK and PG wrote the main manuscript text. JKG, PG and SHA prepared figures. SHA processed the CMIP5 simulations. JKG, PG and AGK designed the study. All authors reviewed and edited the manuscript.

## **Reprints and permissions**

Reprints and permissions information is available at [www.nature.com/reprints](http://www.nature.com/reprints).

## **Competing financial interests**

The authors declare no competing financial interests.

## Figure Captions

**Figure 1. Atmospheric forcings and biospheric forcings.**  $X \rightarrow Y$  represents the fraction of variance of Y explained by X, for the *atmospheric forcing* (atmosphere  $\rightarrow$  biosphere) (**a,c**), and *biospheric forcing* (biosphere  $\rightarrow$  atmosphere) (**b,d**). The signs of the fractions in the top row show whether the atmospheric variable increases (positive) or decreases (negative) the biosphere flux, while in the bottom row they show whether the biosphere increases or decreases the atmospheric response. Oceans and regions where SIF partial correlations are less than 0.1 are shown in white. Pixels without significance are shown in gray (p-value<0.1).

**Fig. 2. Hotspots of terrestrial biosphere-atmosphere feedbacks.** The fraction of biosphere-atmosphere coupling variance explained for the full feedback loop: precipitation  $\rightarrow$  SIF  $\rightarrow$  precipitation (**a**) and PAR  $\rightarrow$  SIF  $\rightarrow$  PAR (**b**). The sign of the fraction shows whether the feedback is positive or negative. Oceans and regions where SIF partial correlations are less than 0.1 are shown in white. Pixels without significance are shown in gray (p-value<0.1).

**Fig. 3. Comparison of observational and Earth System Model results.** Boxplots showing the distributions of significant observational and model results for the fractions of variance explained for the feedbacks of precipitation  $\rightarrow$  biosphere  $\rightarrow$  precipitation (**a**) and PAR  $\rightarrow$  biosphere  $\rightarrow$  PAR (**b**). Boxes are defined by the upper quartile, median and lower quartile of the data while whiskers are defined by the outliers. Only significant pixels are represented (p-value<0.1).

## Methods

### Datasets

Observational remote sensing data is used for SIF, precipitation, and PAR, while quasi-observational reanalysis data is used for temperature. GOME-2, version 2.6<sup>22</sup> (overpass time of 9:30am) is used for SIF, precipitation data is obtained from version 1.2 of GPCP<sup>23</sup>, PAR from CERES<sup>24</sup>, and surface air temperature (1000mb) data from ERA-Interim<sup>25</sup> (see Data availability). While a longer observational data record would allow further insight into interannual variability, we are limited by the satellite data record availability.

There is a certain amount of uncertainty inherent to each product that is described in detail in their data quality summaries. The SIF data is especially noisy (particularly in South America where there are less frequent measurements due to clouds, specifically in the rainforest, and noise from the South Atlantic Anomaly)<sup>22</sup>. Thus, in addition to a standard normalization (described below), SIF data is averaged with the 8 adjacent pixels surrounding the pixel of interest to smooth the remaining noise. On rare pixels, we note that SIF appears to cause an increase in both precipitation and PAR (Figs 1b and d) but this effect is attributed to the use of nine-pixels spatially smoothing of the SIF signal.

The monthly SIF data is calculated from daily measurements (level 2) when the effective cloud fraction is <30%. It should be noted that effective cloud fraction is not equivalent to geometric cloud fraction but is instead based on a Lambertian model that considers cloud reflectance and albedo<sup>44,45,46</sup>. It has been demonstrated that in a typical pixel with a true cloud fraction of 40% that over 80% of the SIF signal can still be retrieved for very thick cloud optical thicknesses (up to 10)<sup>47</sup>. The effective cloud fraction is typically lower than the geometric one.

While cloud filtering could result in a slight bias, it has been shown that altering the effective cloud fraction threshold between 0 and 50 percent only minimally affects the spatial and temporal patterns of SIF<sup>22</sup>. Therefore, we expect minimal bias due to the filtering at the monthly resolution that we consider in our analysis. The one region where the cloud coverage filtering may reduce G-causality detected is in the wet tropics where there is a higher prevalence of clouds. It is possible that the PAR → SIF → PAR feedbacks might be underestimated in this region because of the cloud contamination.

### **SIF-GPP relationship**

This study uses SIF as a proxy for GPP. SIF is mechanistically linked to GPP<sup>9,48</sup>, through both light use efficiency and fAPAR<sup>49</sup>, and has been shown to have a near-linear relationship with GPP at both canopy and ecosystem scales<sup>11,12,50,51,46,52</sup>. While the hourly leaf-level relationship between SIF and GPP has been estimated as curvilinear (SIF continues to increase after the maximum rate of photosynthesis has been reached)<sup>11</sup>, the relationship at larger and longer time scales (e.g., monthly) becomes linear likely due to the effects of averaging across a canopy of leaves representing varying light conditions<sup>11</sup>.

The linearity between SIF and GPP has been observed across biomes using a variety of datasets, including flux tower validation<sup>46,52</sup>. As is shown in Supplementary Fig. 1, SIF correlates strongly with monthly global GPP estimates from Fluxnet-MTE in regions outside of the wet tropics. The SIF-GPP correlation is lower in the wet tropics as the machine learning upscaling approach of the Fluxnet-MTE GPP product has the greatest uncertainty in these regions, as there are few(er) eddy covariance towers there that are used for training<sup>53,54</sup>. Additionally, tropical forest GPP exhibits minimal seasonality<sup>55</sup>, and thus the lower correlation can be attributed to the fitting of noise ( $R^2$  by construction will be small). It has nonetheless been shown that the minimal seasonality in SIF observed in the Amazon

correctly corresponds to the seasonality of carbon dioxide<sup>56</sup> and MODIS near-infrared reflectance related to photosynthesis<sup>55</sup>. As a result, SIF has been used as a proxy for GPP interannual variability<sup>11</sup>.

The linear scaling factor between SIF and GPP varies spatially. Yet, when we normalize the data prior to running the G-Causality, the differing slope values should not impact results since we look at each pixel (location/ecosystem) separately.

### **Conditional MVGC**

We base our analysis on Multivariate Granger causality, using a MVGC MATLAB toolbox<sup>21</sup>, which allows for time and frequency domain MVGC analysis of time series data. The method fits multivariate VAR models to time series. Conditional MVGC compares VAR models with and without (potentially causal) variables. For example, if the addition of past values of precipitation improves the quality of the VAR model prediction for SIF (that uses the autoregressive histories of other variables: SIF, PAR and temperature), then precipitation is considered to have a G-causal influence<sup>57</sup>. If there is no significant information gained (based on an F-test with a null-hypothesis of no G-causality), then the variables are considered not to have a causal link.

Prior to applying the MVGC technique, the data obtained are aggregated to 1-degree by 1-degree monthly data. Monthly data are used to reduce random noise in the original SIF daily data and to achieve consistency with the monthly-aggregated resolution of Coupled Model Intercomparison Project Phase 5 (CMIP5) model data. For each dataset, the long-term mean value is subtracted from each pixel and it is normalized by its long-term standard deviation. After normalization, SIF data is averaged with the 8 adjacent pixels surrounding the pixel of interest to smooth the remaining noise inherent in the SIF data from GOME-2. Single missing monthly values (approximately 4% of the pixels per month) are interpolated

using temporal splines. Prior to performing the normalization and running the MVGC analysis, partial correlations are calculated between non-normalized SIF and atmospheric variables, and if the absolute correlation falls below a value of 0.1, the atmospheric variable is considered non-significant for that pixel and is not included in the analysis. Although results of the analysis are not shown for surface air temperature (temperature at 1000mb), it is used in the analysis, to account for its influence when determining the feedbacks involving precipitation, PAR and SIF. For example, by including temperature in the analysis we guarantee that the G-causality between PAR and SIF is not instead a reflection of the effects of temperature (or related to vapor pressure deficit), which can be correlated with PAR. For all analyses, we use a conservative p-value calculation given the high auto-correlation in the variables of interest, which reduces the degrees of freedom in the number of samples.

Note that we intentionally do not remove the seasonal cycle in pre-processing. Small stochastic amplitude and phase modulations of the seasonality (e.g. large monthly cloud cover or colder than usual temperatures in a particular year) induce non-additive widening of the amplitude and phase spectra so that subtracting the climatology artificially reduces specific frequencies and phases, potentially removing part of the causal signal. This risk is amplified by the relatively short remote sensing record used, which could lead to an imperfect definition of the climatological seasonal cycle. Indeed, where the seasonal signal amplitude and phase have a causal effect we want to capture this (such as the rainfall impact on vegetation green-up and SIF in monsoonal regions). Because the VAR models can capture seasonal periodicity, the MVGC analysis is not affected by the risk of false attribution of causality due to simple lagged seasonality, as is further demonstrates in the examples below.

After normalization of the data and checking that partial correlations between SIF and the other variables fall above 0.1, the Akaike information criterion is calculated and defines the best model order up to the maximum model order, specified as 6 months

442 ('tsdata\_to\_infocrit.m' function in the MVGC MATLAB toolbox). The best actual model  
 443 order used displays the memory of the biosphere-atmosphere interactions (Supplementary  
 444 Fig. 9): model orders of 1 correspond to regions where memory in the system is short and  
 445 causal influence between the atmosphere and biosphere is weak. Using the calculated model  
 446 order, an ordinary least-square regression is used to determine the multivariate-VAR model  
 447 coefficients ('tsdata\_to\_var.m'). The autocovariance function is created  
 448 ('var\_to\_autocov.m'), and from this we calculate the time domain pair-wise conditional  
 449 causalities ('autocov\_to\_pwcgc.m'). To test time-domain significance, we calculate the p-  
 450 values, which are compared to our chosen p-value of less than 0.1 ('mvgc\_pval.m'). An F-  
 451 test with a null-hypothesis of no G-causality is used and only significant pixels are displayed  
 452 in figures. To perform the analysis in the frequency domain and identify subseasonal (<3  
 453 months), seasonal (3 to 12 months) and interannual (>1 year) feedbacks, we calculate the  
 454 spectral-conditional G-causality ('autocov\_to\_spwcgc.m') (Supplementary Fig. 3).

455 We check that the G-causality in the frequency domain integrates to the time domain  
 456 by integrating the frequency results ('smvgc\_to\_mvgc.m') and then subtracting the output  
 457 from the time domain result. Checks are performed throughout the process so that the  
 458 analysis is automatically exited should there be a failed calculation.

459 A sample first order VAR model to explain the variability of SIF is displayed in  
 460 equation 1 with  $A$ ,  $P$ ,  $T$  and  $sig$  representing the VAR coefficient matrix, precipitation,  
 461 temperature, and significance (1 for significant, 0 for insignificant at  $p < 0.1$ ) accordingly.

$$\begin{aligned}
 SIF(t) = & A_{(SIF)} SIF_{(t-1)} + A_{(P \text{ on } SIF)} P_{(t-1)} sig_{(P \text{ on } SIF)} \\
 & + A_{(PAR \text{ on } SIF)} PAR_{(t-1)} sig_{(PAR \text{ on } SIF)} \\
 & + A_{(T \text{ on } SIF)} T_{(t-1)} sig_{(T \text{ on } SIF)} + \varepsilon
 \end{aligned} \tag{1}$$

462 With the addition of the auto-regressive histories of each variable, the VAR model  
 463 captures the original SIF data more accurately. We acknowledge that other factors not  
 464 included in this analysis can affect SIF variability (such as naturally and anthropogenically  
 465 caused disturbances), and is one of the reasons (along with sensor noise) that we cannot  
 466 predict 100% of the variable variance, even with our full VAR model.

### 467 **Synthetic Bootstrap Tests**

468 To demonstrate the effectiveness of this method, we perform several additional tests  
 469 of the conditional MVGC on synthetic data where causal links can be specified. In the first  
 470 three test scenarios PAR and precipitation (P) time series are assumed to be sinusoidal with  
 471 amplitude modulation – AM – and frequency modulation – FM –, as well as additive noise  
 472 (equations 2 and 3). We define two similar test cases except that one has a causal link  
 473 (equation 4) while the other does not (non-causal) (equation 5). We assume that the noise is  
 474 normally distributed (and thus have a white noise/flat spectrum in the frequency domain). To  
 475 test the frequency response, PAR is assumed to have a yearly frequency  
 476  $\omega = 2\pi / (12 \text{ months})$  (equation 2) while precipitation is assumed to have twice-yearly  
 477 frequency  $2\omega$  (i.e. two wet/dry seasons per year) (equation 3).

$$PAR(t) = 100(1 + 0.25A_t^{PAR})\sin\left(\left(1 + \frac{1}{24}F_t^{PAR}\right)\omega t - \pi/2\right) + 25\varepsilon_t^{PAR} \quad (2)$$

$$P(t) = 100(1 + 0.25A_t^P)\sin\left(\left(1 + \frac{1}{24}F_t^P\right)2\omega t - \pi/4\right) + 25\varepsilon_t^P, \quad (3)$$

478 with  $A_t^{PAR}$ ,  $F_t^{PAR}$ ,  $\varepsilon_t^{PAR}$ ,  $A_t^P$ ,  $F_t^P$ ,  $\varepsilon_t^P$  i.i.d. normally distributed with unit variance  $N(0,1)$ .

479 In the *causal case*, SIF is defined as a lagged version of precipitation and radiation  
 480 (with  $t$  in months) (equation 4):

$$SIF = 0.2(1 + 0.25A_t^{SIF})P(t-2) + 0.8(1 + 0.25B_t^{SIF})PAR(t-1) + 25\epsilon_t^{SIF} \quad (4)$$

with  $A_t^{SIF}, B_t^{SIF}, \epsilon_t^{SIF}$  i.i.d. normally distributed with unit variance  $N(0,1)$ . We use 50 years of synthetic data and one realization for the test.

The conditional G-causality finds that only radiation and precipitation are causing SIF and not the converse (Supplementary Fig. 10). In addition, the magnitude of radiation on SIF is four times stronger than the one of precipitation on SIF, as expected based on the time series generated (equation 4).

To emphasize that these results are not spurious, we perform a second, similar test but with a *non-causal* time series (equation 5). This non-causal SIF time series is not induced by PAR nor precipitation. It is statistically similar to the causal scenario, composed of lagged sinusoids with similar frequencies to PAR and precipitation, but without a causal mechanism. For the precipitation and radiation time series we allow for both amplitude and frequency modulations so that both amplitude and phase are stochastic (similar to radiation and precipitation monthly time series).

$$\begin{aligned} SIF = & 20(1 + 0.25a_t^{SIF})\sin\left(\left(1 + \frac{1}{24}b_t^{SIF}\right)2\omega t - \pi/2 - (2/12)2\pi\right) \\ & + 80(1 + 0.25c_t^{SIF})\sin\left(\left(1 + \frac{1}{24}d_t^{SIF}\right)\omega t - \pi/4 - (1/12)2\pi\right) \\ & + 25e_t^{SIF} \end{aligned} \quad (5)$$

The conditional MVGC analysis of this non-causal time series shows no significant G-causality, as expected (Supplementary Fig. 10).

In the third test we bootstrap every month of equations 2-4 across years, clearly destroying the causality in the time series (as the same month from another year is used) while preserving the climatology (and seasonal cycle). As seen in Supplementary Figure 10,

the test again finds no causality in the time series, further confirming the quality of the method and its applicability for our type of time series.

In a fourth and final synthetic data analysis, we test whether we can detect a *causal* full-feedback loop. We repeat the original *causal* test (equation 4), switching the original equation for PAR (equation 2) for one that also includes SIF as a driver (equation 6).

$$PAR = PAR + 0.4 SIF \text{ var}(PAR)/\text{var}(SIF). \quad (6)$$

As expected, in addition to the causality detected previously in the *causal* test of precipitation and PAR on SIF, we also detect significant causality of SIF on PAR (Supplementary Fig. 10).

#### **Observational Bootstrap Test**

To further test the assumption that the observed causation of the biosphere on the atmosphere is not an artifact of the seasonal cycle, we perform a bootstrap analysis with 100-realizations at the global scale. Observational data is sampled by randomly swapping the same months across years for each variable: that is the seasonality is preserved while the causal link from month to month is destroyed. As expected, very few pixels showed any G-causality (Supplementary Fig. 2): only 6.2% of the SIF → precipitation results, and 6.9% of the SIF → PAR results were found to be significant at the 95% confidence level (had more than 5/100 realizations per pixel with significant results based on an F-distribution with a p-value < 0.1). The resulting averaged pair-wise conditional G-causality shows almost no signal, with a peak of less than 0.05 compared to 0.3 for the original dataset (Supplementary Fig. 2). In addition, the resulting geographical patterns reflect mostly random noise. This further emphasizes the physical nature of our assessed causation between the biosphere and the atmosphere.

#### **Vector Autoregressive Models**

The VAR models obtained from the G-causality analysis are used to quantify the fraction of variance in the biosphere explained by the atmosphere and vice versa. We tested for normality and homoscedasticity of the residuals during the VAR fits and excluded pixels that did not meet these criteria (3-6% of pixels depending on the feedback). Using the VAR coefficients generated by the analysis (to account for cross variations), VAR models are created for each atmospheric variable with and without the inclusion of SIF. VAR models are also created for SIF with and without the inclusion of each atmospheric variable. The fractions of observed SIF variance explained by each atmospheric component is computed (equation 7):

$$f_{X \rightarrow SIF} = \frac{\text{var}(SIF_{AR \text{ fit with } X}) - \text{var}(SIF_{AR \text{ fit without } X})}{\text{var}(SIF)} \quad (7)$$

as well as the fraction of each atmospheric variable observed variance explained by SIF (equation 8) (Fig. 1):

$$f_{SIF \rightarrow Y} = \frac{\text{var}(Y_{AR \text{ fit with SIF}}) - \text{var}(Y_{AR \text{ fit without SIF}})}{\text{var}(Y)} \quad (8)$$

These are combined to obtain the full feedback fractions (equation 9) (Fig. 2 and Supplementary Fig. 5):

$$f_{X \rightarrow SIF \rightarrow Y} = \frac{\text{var}(SIF_{AR \text{ fit with } X}) - \text{var}(SIF_{AR \text{ fit without } X})}{\text{var}(SIF)} \times \frac{\text{var}(Y_{AR \text{ fit with SIF}}) - \text{var}(Y_{AR \text{ fit without SIF}})}{\text{var}(Y)} \quad (9)$$

The feedback is defined as positive or negative by taking the VAR model first order coefficients, which is then compared with the VAR model coefficient with the greatest absolute magnitude as further verification. The leading order coefficient of the AR model

could be used in lieu of the first order one but given the rapid decay of the autocorrelation function and the reduced VAR model order (typically less than 2, Supplementary Fig. 9) we use the sign of the first order coefficient. The two estimates of the sign differ in limited regions.

## **CMIP5 Model Simulations**

For the Earth System models from the CMIP5 collection (Supplementary Table 1), the same analysis used for the observational data is applied. Only those models that included GPP data are used. The time period of 1956-2005 is used to obtain statistics that are robust across interannual variability<sup>40</sup>. The true feedback strengths have likely not changed significantly from this earlier, longer time period and the period used for the observational analysis, but we acknowledge that land-use and land-cover changes can affect the feedback metrics (but are also model dependent). One realization of the historical run was used for each model<sup>58</sup>.

VAR models are created based on coefficients calculated in the MVGC analysis for each ESM, and the fraction of variance explained in biosphere-atmosphere coupling from each variable is calculated using equations 5-7.

## **Code availability**

The code used as the basis for the study can be accessed from <http://www.sussex.ac.uk/sackler/mvgc/>.

## **Data availability**

All data supporting the findings of this study are freely available from the following locations:






- GOME-2 SIF: [https://avdc.gsfc.nasa.gov/pub/data/satellite/MetOp/GOME\\_F/](https://avdc.gsfc.nasa.gov/pub/data/satellite/MetOp/GOME_F/)




561       • GPCP precipitation:  
562       <http://iridl.ldeo.columbia.edu/SOURCES/.NASA/.GPCP/.V1DD/.V1p2/>  
563       • CERES PAR: <https://ceres-tool.larc.nasa.gov/ord-tool/jsp/SYN1degSelection.jsp>  
564       • CERES cloud coverage:  
565       <https://ceres-tool.larc.nasa.gov/ord-tool/jsp/ISCCP-D2Selection.jsp>  
566       • ERA-Interim temperature and boundary layer height:  
567       <http://apps.ecmwf.int/datasets/data/interim-full-mnth/levtype=sfc/>  
568       • Fluxnet-MTE surface flux and GPP data:  
569       <https://www.bgc-jena.mpg.de/geodb/projects/Data.php>  
570       • CMIP5 model data: <https://pcmdi.llnl.gov/>  
571       Additional intermediate datasets produced as part of the study can be made available  
572       upon request.


## 573    **Methods References**

- 574    44.    Koelemeijer, R. B. a., Stammes, P., Hovenier, J. W. & de Haan, J. F. A fast method for  
575           retrieval of cloud parameters using oxygen A band measurements from the Global  
576           Ozone Monitoring Experiment. *J. Geophys. Res.* **106**, 3475 (2001).
- 577    45.    Stammes, P. *et al.* Effective cloud fractions from the Ozone Monitoring Instrument:  
578           Theoretical framework and validation. *J. Geophys. Res. Atmos.* **113**, 1–12 (2008).
- 579    46.    Joiner, J. *et al.* The seasonal cycle of satellite chlorophyll fluorescence observations  
580           and its relationship to vegetation phenology and ecosystem atmosphere carbon  
581           exchange. *Remote Sens. Environ.* **152**, 375–391 (2014).
- 582    47.    Joiner, J. *et al.* Filling-in of near-infrared solar lines by terrestrial fluorescence and  
583           other geophysical effects: Simulations and space-based observations from  
584           SCIAMACHY and GOSAT. *Atmos. Meas. Tech.* **5**, 809–829 (2012).
- 585    48.    Lee, J. *et al.* Forest productivity and water stress in Amazonia : observations from  
586           GOSAT chlorophyll fluorescence Forest productivity and water stress in Amazonia :  
587           observations from GOSAT chlorophyll fluorescence. *Proc. R. Soc. B* **280**, 20130171  
588           (2013).
- 589    49.    Duveiller, G. & Cescatti, A. Spatially downscaling sun-induced chlorophyll  
590           fluorescence leads to an improved temporal correlation with gross primary  
591           productivity. *Remote Sens. Environ.* **182**, 72–89 (2016).
- 592    50.    Guanter, L. *et al.* Retrieval and global assessment of terrestrial chlorophyll  
593           fluorescence from GOSAT space measurements. *Remote Sens. Environ.* **121**, 236–251  
594           (2012).
- 595    51.    Guanter, L. *et al.* Global and time-resolved monitoring of crop photosynthesis with

- 596 chlorophyll fluorescence. *Proc. Natl. Acad. Sci. U. S. A.* **111**, E1327-33 (2014).
- 597 52. Yang, X., Tang, J., Mustard, J. F., Lee, J. & Rossini, M. Solar-induced chlorophyll  
598 fluorescence correlates with canopy photosynthesis on diurnal and seasonal scales in a  
599 temperate deciduous forest. **42**, 2977–2987 (2015).
- 600 53. Anav, A. *et al.* Reviews of Geophysics primary production : A review. *Rev. Geophys.*  
601 787–818 (2015).
- 602 54. Jung, M. *et al.* Global patterns of land-atmosphere fluxes of carbon dioxide, latent  
603 heat, and sensible heat derived from eddy covariance, satellite, and meteorological  
604 observations. *J. Geophys. Res.* **116**, G00J07 (2011).
- 605 55. Xu, L. *et al.* Satellite observation of tropical forest seasonality: spatial patterns of  
606 carbon exchange in Amazonia. *Environ. Res. Lett.* **10**, 84005 (2015).
- 607 56. Parazoo, N. C. *et al.* Interpreting seasonal changes in the carbon balance of southern  
608 Amazonia using measurements of XCO<sub>2</sub> and chlorophyll fluorescence from GOSAT.  
609 *Geophys. Res. Lett.* **40**, 2829–2833 (2013).
- 610 57. Granger, C. W. J. Testing for causality. A personal viewpoint. *J. Econ. Dyn. Control* **2**,  
611 329–352 (1980).
- 612 58. Taylor, K. E., Stouffer, R. J. & Meehl, G. A. An overview of CMIP5 and the  
613 experiment design. *Bull. Am. Meteorol. Soc.* **93**, 485–498 (2012).



**a**Precip  $\rightarrow$  SIF  $\rightarrow$  Precip**b**PAR  $\rightarrow$  SIF  $\rightarrow$  PAR

**a**Precip  $\rightarrow$  Biosphere  $\rightarrow$  Precip**b**PAR  $\rightarrow$  Biosphere  $\rightarrow$  PAR